

Best management practices for INVASIVE NONNATIVE PLANT SPECIES in woods in Rose Valley

Invasive nonnative species	Growth form	Hand-pull repeatedly (dispose of plants in trash bags if flowering or fruiting).	Cut or mow repeatedly (or girdle stem bark).	Cut at ground level and paint glyphosate (Roundup™ or Rodeo™) or triclopyr (Garlon™) on cut stem.	Paint penetrating formulation of triclopyr on bark of intact stem.	Spread corn gluten meal or imazipic (Journey™) on soil in early spring.
Amur honeysuckle	shrub	if less than 3' tall	✓	✓		
border privet	shrub	if less than 3' tall	✓	✓		
burningbush	shrub	if less than 3' tall	✓	✓		
English ivy	woody vine	✓	✓	✓		
garlic mustard	biennial herb	✓				
goutweed	perennial herb	✓	✓	Apply glyphosate or triclopyr on foliage in early spring.		
Japanese barberry	shrub	if less than 3' tall	✓	✓		
Japanese honeysuckle	woody vine	✓	✓			
Japanese knotweed	perennial herb	✓	✓	✓		
Japanese stiltgrass	annual grass	✓	✓			✓
lesser celandine	perennial herb	✓		Spray white vinegar on foliage in early spring.		
mile-a-minute	annual vine	✓				✓
multiflora rose	shrub	if less than 3' tall	✓	✓	✓	
Norway maple	tree	if less than 3' tall	✓	✓	✓	
oriental bittersweet	woody vine	if less than 3' tall	✓	✓		